



52<sup>nd</sup> Austrian Mathematical Olympiad  
Regional Competition—Solutions  
25th March 2021

**Problem 1.** Let  $a$  and  $b$  be positive integers and  $c$  be a positive real number satisfying

$$\frac{a+1}{b+c} = \frac{b}{a}.$$

Prove that  $c \geq 1$  holds.

(Karl Czakler)

*Solution.* The following equations are equivalent to the given equation:

$$\begin{aligned} a^2 + a &= b^2 + bc \\ 4a^2 + 4a + 1 &= 4b^2 + 4bc + 1 \\ (2a + 1)^2 &= 4b^2 + 4bc + 1. \end{aligned}$$

Assume to the contrary that  $c < 1$  holds. This yields

$$(2b)^2 = 4b^2 < (2a + 1)^2 = 4b^2 + 4bc + 1 < 4b^2 + 4b + 1 = (2b + 1)^2.$$

This is a contradiction as the square of an integer cannot lie strictly between two consecutive square numbers. Therefore,  $c \geq 1$  holds and, for instance,  $a = b$  yields  $c = 1$  and therefore there is a solution of the equation with  $c \geq 1$ .

(Karl Czakler)  $\square$

**Problem 2.** Let  $ABC$  be an isosceles triangle with  $AC = BC$  and circumcircle  $k$ . The point  $D$  lies on the shorter arc of  $k$  over the chord  $BC$  and is different from  $B$  and  $C$ . Let  $E$  denote the intersection of  $CD$  and  $AB$ .

Prove that the line through  $B$  and  $C$  is a tangent of the circumcircle of the triangle  $BDE$ .

(Karl Czakler)

*Solution.* We denote the center of the circumcircle of the triangle  $BDE$  by  $M$  and  $\angle BAC = \angle CBA$  by  $\alpha$ . Since the quadrilateral  $ABDC$  is cyclic, we obtain  $\angle BDE = \alpha$ . By the inscribed angle theorem,  $\angle BME = 2\alpha$  and thus  $\angle EBM = \angle MEB = 90^\circ - \alpha$ . Therefore,

$$180^\circ = \angle CBA + \angle MBC + \angle EBM = \alpha + \angle MBC + 90^\circ - \alpha = \angle MBC + 90^\circ$$

holds and we get

$$\angle MBC = 90^\circ,$$

completing the proof.

(Karl Czakler)  $\square$

**Problem 3.** The numbers  $1, 2, \dots, 2020$  and  $2021$  are written on a blackboard. The following operation is executed:

Two numbers are chosen, both are erased and replaced by the absolute value of their difference.

This operation is repeated until there is only one number left on the blackboard.

(a) Show that  $2021$  can be the final number on the blackboard.



*Answer.* There are ten triples satisfying the three conditions. They are given by  $(1, 1, 1)$ ,  $(1, 1, 2)$ ,  $(1, 3, 2)$ ,  $(3, 5, 4)$  and their cyclic permutations.

*Solution.* For the sake of readability, we use the notation  $a \mid b + c$  instead of  $a \mid (b + c)$  throughout the proof.

Without loss of generality, let  $x$  be the smallest of the three numbers (or one of the smallest), i.e.  $x \leq y$  and  $x \leq z$ . From  $z \mid x + 1$  we obtain  $x \leq z \leq x + 1$ . Thus we have to consider two cases.

- Case 1. Let  $z = x$ . Then  $z = x \mid x + 1$  leads to  $x = z = 1$  and  $y \mid z + 1 = 2$ . Therefore  $y = 1$  or  $y = 2$ , and we get the two solutions  $(1, 1, 1)$  and  $(1, 2, 1)$ .
- Case 2. Let  $z = x + 1$ . Then the two conditions  $x \mid y + 1$  and  $y \mid x + 2$  must be fulfilled. In particular, we obtain  $x \leq y + 1$  and  $y \leq x + 2$ . This yields  $x - 1 \leq y \leq x + 2$  and we have to examine the following cases for  $y$ .
  - Case 2a. Let  $0 < y = x$ . The conditions  $x \mid x + 1$  and  $x \mid x + 2$  can only hold simultaneously for  $x = 1$ , giving the solution  $(1, 1, 2)$ .
  - Case 2b. Let  $y = x + 1$ . Then the two conditions are  $x \mid x + 2$  and  $x + 1 \mid x + 2$ . They cannot hold simultaneously.
  - Case 2c. Let  $y = x + 2$ . The condition  $y = x + 2 \mid x + 2 = z + 1$  is trivially fulfilled. The requirement  $x \mid y + 1 = x + 3$  can only hold for  $x \mid 3$ . And, indeed, for either  $x = 1$  or  $x = 3$  the condition is fulfilled and we obtain the solutions  $(1, 3, 2)$  and  $(3, 5, 4)$ .

Summing up, the triples  $(1, 1, 1)$ ,  $(1, 2, 1)$ ,  $(1, 1, 2)$ ,  $(1, 3, 2)$  and  $(3, 5, 4)$  fulfill all three conditions.

As each of the three numbers can be the minimum, every cyclic permutation of these triples is a solution as well.

*(Lukas Donner)*  $\square$